

## SD and k-SD Prime Cordial graphs

A. Delman<sup>1</sup>, S. Koilraj<sup>2</sup> and P. Lawrence Rozario Raj<sup>3</sup>

<sup>2,3</sup>PG and Research Department of Mathematics, St. Joseph's College  
Tiruchirappalli -620002, Tamilnadu, India

E-mail : [skoilraj@yahoo.com](mailto:skoilraj@yahoo.com) , [lawraj2006@yahoo.co.in](mailto:lawraj2006@yahoo.co.in)

<sup>1</sup>PG and Research Department of Mathematics, St. Joseph's College  
Tiruchirappalli -620002, Tamilnadu, India. E-mail: [delmanmaths@gmail.com](mailto:delmanmaths@gmail.com)

Received 17 March 2018; accepted 21 April 2018

**Abstract.** In this paper, we investigate the SD-Prime cordial labeling of  $Pl_n$  graph and k-SD-Prime cordial labeling of  $(P_n \odot K_1) \cup K_{1,n,n}$  and  $P_n \cup K_{1,n,n}$ .

**Keywords:** SD-Prime cordial labeling, k-SD-Prime cordial labeling, k-SD-Prime cordial graph.

**AMS Mathematics Subject Classification (2010):** 05C78

### 1. Introduction

By a graph, we mean a finite, undirected graph without loops and multiple edges, for terms not defined here, we refer to Harary [5]. For standard terminology and notations related to number theory we refer to Burton [2] and graph labeling, we refer to Gallian [4]. The notion of prime labeling for graphs originated with Roger Entringer and was introduced in a paper by Tout et al. [12] in the early 1980s and since then it is an active field of research for many scholars. In [13], Vaidya et al. introduced the concept of k-prime labeling of graph. Sundaram et al. introduced the notion of prime cordial labeling in [11]. The concept of neighborhood-prime labeling of graph was introduced by Patel et al. [10]. Lawrence et al. introduced the notation of k-neighborhood-prime labeling of graph in [8]. Lau et al was introduced a variant of prime graph labeling of graph in [6]. In [7], Lau et al. introduced SD-prime cordial labeling and they discussed SD-prime cordial labeling for some standard graphs. In [9], Lourdusamy et al. investigated some new construction of SD-prime cordial graph. In [3], Delman et.al., introduced the concept of k-SD-prime cordial labeling of graph and discussed k-SD-prime cordial labeling of some standard graphs. In [1], Babujee defined a class of planar graph as graph obtained by removing certain edges from the corresponding complete graph. The class of planar graph so obtained is denoted by  $Pl_n$ . Here we discuss the SD-Prime cordial labeling of  $Pl_n$  graph, for  $n \geq 3$  and k-SD-Prime cordial labeling of  $(P_n \odot K_1) \cup K_{1,n,n}$ , for  $n \geq 2$  and  $P_n \cup K_{1,n,n}$ , for  $n \geq 2$ .

### 2. Basic definitions

**Definition 2.1.** A complete bipartite graph  $K_{1,n}$  is called a star and it has  $n+1$  vertices and  $n$  edges.  $K_{1,n,n}$  is the graph obtained by the subdivision of the edges of the star  $K_{1,n}$ .

A. Delman, S. Koilraj and P. Lawrence Rozario Raj

**Definition 2.2.** Let  $K_n$  be the complete graph on  $n$  vertices  $V_n = \{1, 2, \dots, n\}$ . The class of graphs  $Pl_n$  has the vertex set  $V_n$  and the edge set

$$E_n = E(K_n) \setminus \{(k, l) : 3 \leq k \leq n-2, k+2 \leq l \leq n\}.$$

**Definition 2.3.** Comb is a graph obtained by joining a single pendant edge to each vertex of a path. In other words  $P_n \odot K_1$  is a comb graph.

**Definition 2.4.** Let  $G = (V, E)$  be a graph with  $n$  vertices. A function  $f : V(G) \rightarrow \{1, 2, 3, \dots, n\}$  is said to be a prime labeling, if it is bijective and for every pair of adjacent vertices  $u$  and  $v$ ,  $\gcd(f(u), f(v)) = 1$ . A graph which admits prime labeling is called a prime graph.

**Definition 2.5.** A  $k$ -prime labeling of a graph  $G$  is an injective function  $f : V \rightarrow \{k, k+1, \dots, k+|V|-1\}$  for some positive integer  $k$  that induces a function  $f^+ : E(G) \rightarrow \mathbb{N}$  of the edges of  $G$  defined by  $f^+(uv) = \gcd(f(u), f(v))$ ,  $\forall e = uv \in E(G)$  such that  $\gcd(f(u), f(v)) = 1$ ,  $\forall e = uv \in E(G)$ . The graph which admits a  $k$ -prime labeling is called a  $k$ -prime graph.

**Definition 2.6.** Let  $G = (V, E)$  be a graph with  $n$  vertices. A bijective function  $f : V(G) \rightarrow \{1, 2, 3, \dots, n\}$  is said to be a neighborhood-prime labeling, if for every vertex  $v \in V(G)$  with  $\deg(v) > 1$ ,  $\gcd \{f(u) : u \in N(v)\} = 1$ . A graph which admits neighborhood-prime labeling is called a neighborhood-prime graph.

**Definition 2.7.** Let  $G = (V(G), E(G))$  be a graph with  $n$  vertices. A bijective function  $f : V(G) \rightarrow \{k, k+1, \dots, k+n-1\}$  is said to be a  $k$ -neighborhood-prime labeling, if for every vertex  $v \in V(G)$  with  $\deg(v) > 1$ ,  $\gcd \{f(u) : u \in N(v)\} = 1$ . A graph which admits  $k$ -neighborhood-prime labeling is called a  $k$ -neighborhood-prime graph.

**Definition 2.8.** Given a bijection  $f : V(G) \rightarrow \{1, 2, \dots, |n|\}$ , we associate 2 integers  $S = f(u) + f(v)$  and  $D = |f(u) - f(v)|$  with every edge  $uv$  in  $E$ . The labeling  $f$  induces an edge labeling  $f' : E(G) \rightarrow \{0, 1\}$  such that for any edge  $uv$  in  $G$ ,  $f'(uv) = 1$  if  $\gcd(S, D) = 1$  and 0 otherwise. We say  $f$  is  $SD$ -prime labeling if  $f'(uv) = 1$  for all  $uv \in E(G)$ . Moreover,  $G$  is  $SD$ -prime if it admits  $SD$ -prime labeling.

**Definition 2.9.** Given a bijection  $f : V(G) \rightarrow \{1, 2, \dots, |V(G)|\}$ , we associate two integers  $S = f(u) + f(v)$  and  $D = |f(u) - f(v)|$  with every edge  $uv$  in  $E(G)$ . The labeling  $f$  induces an edge labeling  $f^* : E(G) \rightarrow \{0, 1\}$  such that for any edge  $uv$  in  $E(G)$ ,  $f^*(uv) = 1$  if  $\gcd(S, D) = 1$  and 0 otherwise. Let  $e_{f^*}(i)$  be the number of edges labeled with  $i \in \{0, 1\}$ . We say  $f$  is  $SD$ -prime cordial labeling if  $|e_{f^*}(0) - e_{f^*}(1)| \leq 1$ . Moreover  $G$  is  $SD$ -prime cordial if it admits  $SD$ -prime cordial labeling.

### 3. Main theorems

**Theorem 3.1.**  $Pl_n$  is a  $SD$ -prime cordial graph, for  $n \geq 3$ .

**Proof:** Let  $v_1, v_2, \dots, v_n$  be the vertices and  $e_1, e_2, \dots, e_{3n-6}$  be the edges of  $Pl_n$ , where  $e_i = v_i v_{i+1}$  for  $1 \leq i \leq n-3$ ,  $e_{i+n-3} = v_{n-1} v_i$  for  $1 \leq i \leq n-2$ ,  $e_{i+2n-5} = v_n v_i$  for  $1 \leq i \leq n-2$  and  $e_{3n-6} = v_{n-1} v_n$ .

Let  $G = Pl_n$ . Then  $|V(G)| = n$  and  $|E(G)| = 3n-6$ .

### SD and k-SD Prime Cordial graphs

Define  $f : V(G) \rightarrow \{1, 2, \dots, n\}$  as follows:

**Case 1:**  $n \equiv 1, 3 \pmod{4}$ .

$$g(v_i) = \begin{cases} i+2 & \text{if } i \equiv 0, 1 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ i+3 & \text{if } i \equiv 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ i+1 & \text{if } i \equiv 3 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ 2 & \text{if } i = n-1 \\ 1 & \text{if } i = n \end{cases}$$

Then induced edge labels are

$$g^*(e_{2i-1}) = 0, \text{ for } 1 \leq i \leq \frac{n-3}{2}$$

$$g^*(e_{2i}) = 1, \text{ for } 1 \leq i \leq \frac{n-3}{2}$$

$$g^*(e_{n-3+i}) = \begin{cases} 1 & \text{if } i \equiv 1, 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ 0 & \text{if } i \equiv 0, 3 \pmod{4} \text{ and } 1 \leq i \leq n-2 \end{cases}$$

$$g^*(e_{2n-5+i}) = \begin{cases} 0 & \text{if } i \equiv 1, 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ 1 & \text{if } i \equiv 0, 3 \pmod{4} \text{ and } 1 \leq i \leq n-2 \end{cases}$$

$$g^*(e_{3n-6}) = 1$$

In view of the above defined labeling pattern, we have  $e_{f^*}(0)+1 = e_{f^*}(1) = \frac{3n-5}{2}$  and

$$|e_{f^*}(0) - e_{f^*}(1)| \leq 1.$$

Therefore the  $Pl_n$  is a SD-prime cordial graph, for  $n \equiv 1, 3 \pmod{4}$ .

**Case 2:**  $n \equiv 0 \pmod{4}$ .

$$g(v_i) = \begin{cases} i+2 & \text{if } i \equiv 0, 1 \pmod{4} \text{ and } 1 \leq i \leq n-4 \\ i+3 & \text{if } i \equiv 2 \pmod{4} \text{ and } 1 \leq i \leq n-4 \\ i+1 & \text{if } i \equiv 3 \pmod{4} \text{ and } 1 \leq i \leq n-4 \\ n-1 & \text{if } i = n-3 \\ n & \text{if } i = n-2 \\ 2 & \text{if } i = n-1 \\ 1 & \text{if } i = n \end{cases}$$

Then induced edge labels are

$$g^*(e_{2i-1}) = 0, \quad \text{for } 1 \leq i \leq \frac{n-4}{2}$$

$$g^*(e_{2i}) = 1, \quad \text{for } 1 \leq i \leq \frac{n-6}{2}$$

$$g^*(e_{n-4}) = 0,$$

$$g^*(e_{n-3}) = 1,$$

$$g^*(e_{n-3+i}) = \begin{cases} 1 & \text{if } i \equiv 1, 2 \pmod{4} \text{ and } 1 \leq i \leq n-4 \\ 0 & \text{if } i \equiv 0, 3 \pmod{4} \text{ and } 1 \leq i \leq n-4 \end{cases}$$

$$g^*(e_{2n-6}) = 1,$$

$$g^*(e_{2n-5}) = 0,$$

A. Delman, S. Koilraj and P. Lawrence Rozario Raj

$$g^*(e_{2n-5+i}) = \begin{cases} 0 & \text{if } i \equiv 1, 2 \pmod{4} \text{ and } 1 \leq i \leq n-4 \\ 1 & \text{if } i \equiv 0, 3 \pmod{4} \text{ and } 1 \leq i \leq n-4 \end{cases}$$

$$g^*(e_{3n-8}) = 0,$$

$$g^*(e_{3n-7}) = 1,$$

$$g^*(e_{3n-6}) = 1$$

In view of the above defined labeling pattern, we have  $e_{f^*}(0) = e_{f^*}(1) = \frac{3n-6}{2}$  and  $|e_{f^*}(0) - e_{f^*}(1)| \leq 1$ .

Therefore the  $Pl_n$  is a SD-prime cordial graph,  $n \equiv 0 \pmod{4}$ .

**Case 3:**  $n \equiv 2 \pmod{4}$ .

$$g(v_i) = \begin{cases} i+2 & \text{if } i \equiv 0, 1 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ i+3 & \text{if } i \equiv 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ i+1 & \text{if } i \equiv 3 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ 2 & \text{if } i = n-1 \\ 1 & \text{if } i = n \end{cases}$$

Then induced edge labels are

$$g^*(e_{2i-1}) = 0, \quad \text{for } 1 \leq i \leq \frac{n-3}{2}$$

$$g^*(e_{2i}) = 1, \quad \text{for } 1 \leq i \leq \frac{n-3}{2}$$

$$g^*(e_{n-3+i}) = \begin{cases} 1 & \text{if } i \equiv 1, 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ 0 & \text{if } i \equiv 0, 3 \pmod{4} \text{ and } 1 \leq i \leq n-2 \end{cases}$$

$$g^*(e_{2n-5+i}) = \begin{cases} 0 & \text{if } i \equiv 1, 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ 1 & \text{if } i \equiv 0, 3 \pmod{4} \text{ and } 1 \leq i \leq n-2 \end{cases}$$

$$g^*(e_{3n-6}) = 1$$

In view of the above defined labeling pattern, we have  $e_{f^*}(0) = e_{f^*}(1) = \frac{3n-6}{2}$  and  $|e_{f^*}(0) - e_{f^*}(1)| \leq 1$ .

Therefore the  $Pl_n$  is a SD-prime cordial graph, for  $n \equiv 2 \pmod{4}$ .

Therefore the  $Pl_n$  is a SD-prime cordial graph,  $n \geq 3$ .

**Theorem 3.2:** The disconnected graph  $(P_n \odot K_1) \cup K_{1,m,m}$  is k-SD-prime cordial graph, for  $n, m \geq 2$ .

**Proof:** Let  $P_n \odot K_1$  be a comb graph. Let  $v_1, v_2, \dots, v_{2n}$  be the vertices and  $e_1, e_2, \dots, e_{2n-1}$  be the edges of  $P_n \odot K_1$ . Let  $u, u_1, u_2, \dots, u_{2m}$  be the vertices and  $s_1, s_2, \dots, s_{2m}$  be the edges of  $K_{1,m,m}$ .

Let  $G$  be the disconnected graph  $(P_n \odot K_1) \cup K_{1,m,m}$ .

Then  $|V(G)| = 2n+2m+1$  and  $|E(G)| = 2n+2m-1$ .

Define  $g : V(G) \rightarrow \{k, k+1, \dots, k+2n+2m\}$  as follows:

$$g(v_i) = \begin{cases} k+2i-2, & \text{if } 1 \leq i \leq n \\ k+2i-2n-1, & \text{if } n+1 \leq i \leq 2n \end{cases}$$

$$g(u) = k+2n$$

SD and k-SD Prime Cordial graphs

$$g(u_i) = \begin{cases} k + 2n + 2i, & \text{if } 1 \leq i \leq m \\ k + 2n - 2m + 2i + 1, & \text{if } m + 1 \leq i \leq 2m \end{cases}$$

Then induced edge labels are

$$g^*(e_i) = \begin{cases} 0, & \text{if } 1 \leq i \leq n-1 \\ 1, & \text{if } n \leq i \leq 2n-1 \end{cases}$$

$$g^*(s_i) = \begin{cases} 0, & \text{if } 1 \leq i \leq m \\ 1, & \text{if } m + 1 \leq i \leq 2m \end{cases}$$

In view of the above defined labeling pattern, we have  $e_{f^*}(0) + 1 = e_{f^*}(1) = n + m$  and  $|e_{f^*}(0) - e_{f^*}(1)| \leq 1$ .

Therefore the disconnected graph  $(P_n \odot K_1) \cup K_{1,m,m}$  is k-SD-prime cordial graph, for  $n, m \geq 2$ .

**Theorem 3.3.** The disconnected graph  $P_n \cup K_{1,m,m}$  is k-SD-prime cordial graph, for  $n, m \geq 2$ .

**Proof:** Let  $P_n$  be a path graph. Let  $v_1, v_2, \dots, v_n$  be the vertices and  $e_1, e_2, \dots, e_{n-1}$  be the edges of  $P_n$ . Let  $u, u_1, u_2, \dots, u_{2m}$  be the vertices and  $s_1, s_2, \dots, s_{2m}$  be the edges of  $K_{1,m,m}$ .

Let  $G$  be the disconnected graph  $P_n \cup K_{1,m,m}$ .

Then  $|V(G)| = n + 2m + 1$  and  $|E(G)| = n + 2m - 1$ .

Define  $g : V(G) \rightarrow \{k, k+1, \dots, k+n+2m\}$  as follows:

**Case 1:**  $n \equiv 1, 3 \pmod{4}$ .

$$g(v_i) = \begin{cases} k + i - 1 & \text{if } i \equiv 0, 1 \pmod{4} \text{ and } 1 \leq i \leq n \\ k + i & \text{if } i \equiv 2 \pmod{4} \text{ and } 1 \leq i \leq n \\ k + i - 2 & \text{if } i \equiv 3 \pmod{4} \text{ and } 1 \leq i \leq n \end{cases}$$

$$g(u) = k + n$$

$$g(u_i) = \begin{cases} k + n + 2i, & \text{if } 1 \leq i \leq m \\ k + n - 2m + 2i - 1, & \text{if } m + 1 \leq i \leq 2m \end{cases}$$

Then induced edge labels are

$$g^*(e_{2i-1}) = 0, \quad \text{if } 1 \leq i \leq \frac{n-1}{2}$$

$$g^*(e_{2i}) = 1, \quad \text{if } 1 \leq i \leq \frac{n-1}{2}$$

$$g^*(s_i) = \begin{cases} 0, & \text{if } 1 \leq i \leq m \\ 1, & \text{if } m + 1 \leq i \leq 2m \end{cases}$$

In view of the above defined labeling pattern, we have

$$e_{f^*}(0) = e_{f^*}(1) = \frac{n + 2m - 1}{2} \text{ and } |e_{f^*}(0) - e_{f^*}(1)| \leq 1.$$

Therefore  $P_n \cup K_{1,m,m}$  is k-SD-prime cordial graph, for  $n \equiv 0, 1, 3 \pmod{4}$ .

**Case 2:**  $n \equiv 0 \pmod{4}$ .

A. Delman, S. Koilraj and P. Lawrence Rozario Raj

$$g(v_i) = \begin{cases} k+i-1 & \text{if } i \equiv 0,1 \pmod{4} \text{ and } 1 \leq i \leq n \\ k+i & \text{if } i \equiv 2 \pmod{4} \text{ and } 1 \leq i \leq n \\ k+i-2 & \text{if } i \equiv 3 \pmod{4} \text{ and } 1 \leq i \leq n \end{cases}$$

$$g(u) = k+n$$

$$g(u_i) = \begin{cases} k+n+2i, & \text{if } 1 \leq i \leq m \\ k+n-2m+2i-1, & \text{if } m+1 \leq i \leq 2m \end{cases}$$

Then induced edge labels are

$$g^*(e_{2i-1}) = 0, \quad \text{if } 1 \leq i \leq \frac{n}{2}$$

$$g^*(e_{2i}) = 1, \quad \text{if } 1 \leq i \leq \frac{n-2}{2}$$

$$g^*(s_i) = \begin{cases} 0, & \text{if } 1 \leq i \leq m \\ 1, & \text{if } m+1 \leq i \leq 2m \end{cases}$$

In view of the above defined labeling pattern, we have  $e_{f^*}(0) = e_{f^*}(1)+1 = \frac{n+2m}{2}$  and

$$|e_{f^*}(0) - e_{f^*}(1)| \leq 1.$$

Therefore  $P_n \cup K_{1,m,m}$  is  $k$ -SD-prime cordial graph, for  $n \equiv 0 \pmod{4}$ .

**Case 3:**  $n \equiv 2 \pmod{4}$ .

$$g(v_i) = \begin{cases} k+i-1 & \text{if } i \equiv 0,1 \pmod{4} \text{ and } 1 \leq i \leq n-1 \\ k+i & \text{if } i \equiv 2 \pmod{4} \text{ and } 1 \leq i \leq n-1 \\ k+i-2 & \text{if } i \equiv 3 \pmod{4} \text{ and } 1 \leq i \leq n-1 \\ k+n-1 & \text{if } i = n \end{cases}$$

$$f(u) = k+n$$

$$g(u_i) = \begin{cases} k+n+2i, & \text{if } 1 \leq i \leq m \\ k+n-2m+2i-1, & \text{if } m+1 \leq i \leq 2m \end{cases}$$

Then induced edge labels are

$$f^*(e_{2i-1}) = 0 \quad \text{if } 1 \leq i \leq \frac{n-2}{2}$$

$$f^*(e_{2i}) = 1 \quad \text{if } 1 \leq i \leq \frac{n-2}{2}$$

$$f^*(e_{2n-1}) = 1$$

$$g^*(s_i) = \begin{cases} 0, & \text{if } 1 \leq i \leq m \\ 1, & \text{if } m+1 \leq i \leq 2m \end{cases}$$

In view of the above defined labeling pattern, we have  $e_{f^*}(0)+1 = e_{f^*}(1) = \frac{n+2m}{2}$  and

$$|e_{f^*}(0) - e_{f^*}(1)| \leq 1.$$

Therefore  $P_n \cup K_{1,n,n}$  is  $k$ -SD-prime cordial graph, for  $n \equiv 2 \pmod{4}$ .

Hence the disconnected graph  $P_n \cup K_{1,n,n}$  is  $k$ -SD-prime cordial graph, for  $n, m \geq 2$ .

#### 4. Conclusions

In this paper, we presented the SD-Prime cordial labeling of  $Pl_n$  graph, for  $n \geq 3$  and k-SD-Prime cordial labeling of  $(P_n \odot K_1) \cup K_{1,n,n}$ , for  $n \geq 2$  and  $P_n \cup K_{1,n,n}$ , for  $n \geq 2$ .

#### REFERENCES

1. J.Baskar Babujee, Planar graphs with maximum edges-antimagic property, *The Mathematics Education*, 37(4) (2003) 194-198.
2. D.M.Burton, *Elementary Number Theory*, Second Edition, Wm. C. Brown Company Publishers, (1980).
3. A.Delman, S.Koilraj and P.Lawrence Rozario Raj, k-SD-Prime Cordial Labeling of graphs, communicated.
4. J.A.Gallian, A dynamic survey of graph labeling, *The Electronic Journal of Combinatorics*, 16, #DS6, )2016).
5. F.Harary, *Graph theory*, Addison Wesley, Reading, Massachusetts, (1972).
6. G.C.Lau and W.C.Shiu, On SD-prime labeling of graphs, *Utilitas Math.*, accepted.
7. G.C.Lau, H.H.Chu, N.Suhadak, F.Y.Foo and H.K.Ng, On SD-prime cordial graphs. *Int. J. Pure Appl. Math.*, 106(4) (2016) 1017-1028.
8. P.Lawrence Rozario Raj and Sr.Jincy Joseph, k-Neighborhood-prime labeling of graphs, *International Journal of Innovative Science, Engineering & Technology*, 3(8) (2016) 288-292.
9. A.Lourdusamy and F.Patrick, New construction on SD-prime cordial labeling, In: Arumugam S., Bagga J., Beineke L., Panda B. (eds) *Theoretical Computer Science and Discrete Mathematics, ICTCSDM 2016, Lecture Notes in Computer Science, LNCS 10398*, Springer, Cham, pp. 134-143, (2017).
10. S.K.Patel and N.P.Shrimali, Neighborhood-prime labeling, *International Journal of Mathematics and Soft Computing*, 5(2) (2015) 135-143.
11. M.Sundaram, R.Ponraj and S.Somasundaram, Prime cordial labeling of graphs, *J. Indian Acad. Math.*, 27 (2005) 373-390.
12. A.Tout, A.N.Dabboucy and K.Howalla, Prime labeling of graphs, *Nat. Acad. Sci. Letters*, 11 (1982) 365-368.
13. S.K.Vaidya and U.M.Prajapati, Some results on prime and k-prime labeling, *Journal of Mathematics Research*, 3(1) (2011) 66-75.